

Dosing for Infants and Children



From Your Healthcare Professional

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.

If possible, use weight to dose; otherwise, use age.

mL = milliliter



Infants' TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL)

Available in:



Use product only as directed.



Children's TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL)

Available in:



Use product only as directed.

| WEIGHT | AGE | Infants' TYLENOL® | Children's TYLENOL® |
|--------------|--------------|-------------------|---------------------|
| 6-11 POUNDS | 0-3 MONTHS | 1.25 mL | — |
| 12-17 POUNDS | 4-11 MONTHS | 2.5 mL | — |
| 18-23 POUNDS | 12-23 MONTHS | 3.75 mL | — |
| 24-35 POUNDS | 2-3 YEARS | 5 mL | 5 mL |
| 36-47 POUNDS | 4-5 YEARS | — | 7.5 mL |
| 48-59 POUNDS | 6-8 YEARS | — | 10 mL |
| 60-71 POUNDS | 9-10 YEARS | — | 12.5 mL |
| 72-95 POUNDS | 11 YEARS | — | 15 mL |

Kids' Wellness Tracker



This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App StoreSM and the Google Play StoreTM



Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _____

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all TYLENOL® products.
- Repeat dose every 4 hours while symptoms last.
- Do NOT give more than 5 doses in 24 hours.
- Do NOT use with any other product containing acetaminophen.
- Use only the dosing device (syringe or dosing cup) that came with the product.
- **All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL).**